



Drug Information

RITONAVIR (Norvir)

Last updated September 2003

Why am I given RITONAVIR as my medication?

- RITONAVIR (also called Norvir) is a drug used in combination with other medications to treat HIV (Human Immunodeficiency Virus) infection.
- RITONAVIR blocks a protein called “Protease”. It belongs to a class of drugs called Protease Inhibitors (PI). The HIV virus needs this protein to reproduce itself, so by blocking it the drug helps slow down HIV disease.
- Taking this medication can reduce the amount of virus in your body. It can also reduce your chance of getting sick from AIDS related illnesses, help you stay healthy longer or get your health back. It may also reduce the damage to your immune system.

How do I take this medication?

- RITONAVIR comes as a 100 mg capsule and 80mg/cc liquid. The usual dose of RITONAVIR is 600 mg (6 capsules or 7.5 c.c. of the liquid) twice daily.
- NORVIR should be taken with food.
- To reduce side effects, RITONAVIR should be started at a lower dose and increased gradually over 2 weeks (for example, 300 mg twice daily for 3 days, then 400 mg twice daily for 3 days, then 500 mg twice daily for 3 days, then 600 mg twice daily).
- RITONAVIR may be used at a lower dose if it is combined with other protease inhibitor.
- The oral solution can be taken alone or with other fluids to improve its taste. You can mix the RITONAVIR liquid with a cup of chocolate milk, Ensure, Advera or ice cream. Mix only one dose at a time. After mixing, the liquid should be taken within an hour.
- RITONAVIR should be kept at room temperature away from direct sunlight. RITONAVIR capsules can be left at room temperature for 30 days. Store extra amounts of RITONAVIR in the refrigerator.
- Keep it out of reach of children.

What if I forget to take a dose?

- Take the dose you missed as soon as possible. However, if it is within 2 hours of your next dose, just continue with your regular schedule. Do not double the dose.
- Recent studies have found that for the anti-HIV medications to work, all the medications need to be taken regularly and consistently. Missing or skipping dose of your medication may make it lose its effectiveness as the virus can change and become resistant to the medication.



What are the side effects of RITONAVIR?

- The most common side effect of NORVIR include:
 - stomach upset
 - nausea and vomiting
 - diarrhea
 - loss of appetite, taste changes
 - dizziness, headache
 - fatigue
 - tingling and numbness in hands, feet or around the lips.
- Longer term side effects of RITONAVIR can include:
 - increase in the sugar and fat (cholesterol, triglyceride) levels in your blood
 - abnormal body fat distribution (increase in waist and breast size and thinning of the face, arms and legs).
- Consult your doctor or pharmacist if you have these side effects. Do not stop the medication or change the dose before you talk to them.

Can I take RITONAVIR with other medications?

- RITONAVIR can interact with other drugs. It is important that you tell your doctor and pharmacist about all the prescription and non-prescription medications (including vitamins and herbs) you are taking.
- RITONAVIR should not be taken with Halcion (triazolam), Versed (midazolam), Hismanal (astemizole), Seldane (terfenadine), Prepulsid (cisapride), Xanax (alprazolam), Cordarone (amiodarone), Tranxene (clorazepate), Clorzaryl (Clozapine), Valium (diazepam), Enkaid (encainide), Prosom (estazolam), Tambocor (flecainide), Dalmane (flurazepam), Demerol (meperidine), Feldene (piroxicam), Rythmol (propafenone), Darvon (propoxyphene), Quinidine, and Mycobutin (Rifabutin).
- RITONAVIR can decrease the effectiveness of birth control pills. An alternative method for birth control such as latex condoms, should be used if you are on RITONAVIR.
- RITONAVIR may reduce the effectiveness of Methadone. Your Methadone dosage may need to be adjusted if you are on RITONAVIR.

Can I take RITONAVIR with alcohol or street drugs?

- In general, it is advisable to avoid excessive amount of alcohol while you are taking anti-HIV medications. Alcohol may interact with some of your medications. Do not skip a dose of your medication because you want a drink.
- RITONAVIR may also interact with other street drugs (e.g. ecstasy), consult your doctor and pharmacist if you are using street drugs so they can advise you with the necessary precautions.

Can I take RITONAVIR if I am pregnant or breastfeeding?

- If you are pregnant and wish to take RITONAVIR, please consult your doctor.
- Since the HIV virus can be transmitted through breast milk, breast-feeding is not recom-



mended in HIV positive women.

What other precautions do I need to know when taking RITONAVIR?

- You should keep your appointments with your physician for blood tests to check your liver and kidney function, blood sugar, cholesterol and triglyceride regularly.
- Make sure you have a continuous supply of the medication.
- RITONAVIR does not kill the virus or cure AIDS. It also does not prevent the transmission of HIV, so please remember to always take precautions if you are having sex (e.g. use latex condoms) or using drugs (use clean syringes).

Copyright @ACAS 2001 & 2003. This fact sheet is produced by ACAS (Asian Community AIDS Services) and is available in English, Chinese (Traditional and Simplified), Tagalog and Vietnamese. Funding for this project is provided by the Ontario HIV Treatment Network (OHTN) and Health Canada. Copies can be downloaded at : www.acas.org/treatment. ACAS provides comprehensive support and case management services for Asian people living with HIV/AIDS; and HIV/AIDS prevention education to the east and southeast Asian Canadian populations in the Greater Toronto Area. ACAS is located at 33 Isabella Street, Suite 107, Toronto, Ontario M4Y 2P7. Tel: (416)-963-4300, Fax: (416)-963-4371 Email: support@acas.org

Legal Disclaimer: While we make every attempt to ensure the accuracy and reliability of information contained in this website/ fact sheet, the information provided here are designed for reference purposes only. These information should not be relied upon as a substitute for medical advice from a qualified professional health care provider and should not be used for diagnosing or treating a condition or illness. Please consult a physician if you have any concerns about your health, treatment regimen and questions related to HIV/AIDS. ACAS, its employees and board members will not be responsible for any loss or harm, however arising, from the use of, or reliance on this information.