



Drug Information

SAQUINAVIR (Invirase, Fortovase)

Last updated September 2003

Why am I given SAQUINAVIR as my medication?

- SAQUINAVIR (Invirase, Fortovase) is a drug used in combination with other medications to treat HIV (Human Immunodeficiency Virus) infection.
- SAQUINAVIR blocks a protein called “Protease”. It belongs to a class of drugs called Protease Inhibitors (PI). The HIV virus needs this protein to reproduce itself, so by blocking it the drug helps slow down HIV disease.
- Taking this medication can reduce the amount of virus in your body. It can also reduce your chance of getting sick from AIDS related illnesses, help you stay healthy longer or get your health back. It may also reduce the damage to your immune system.

How do I take this medication?

- SAQUINAVIR comes in 2 different forms:
 - INVIRASE, which is a 200 mg green and yellow capsule
 - FORTOVASE, which is a 200 mg beige color soft gel capsule.
- The number of pills you need to take depend on which form of the drug you are taking. Both INVIRASE and FORTOVASE should be taken with food or within two hours after a meal.
- Swallow the capsules whole, do not chew or crush them.
- These days, Saquinavir is mostly used in combination with another Protease Inhibitor, such as Ritonavir (Norvir).
- Don't put the medicine in the bathroom or kitchen, as moisture may cause the medicine to lose its effectiveness.
 - INVIRASE capsules should be kept at room temperature in a dry place.
 - FORTOVASE capsules may be stored at room temperature for up to 3 months.
- Keep it out of reach of children.

What if I forget to take a dose?

- Take the dose you missed as soon as possible. However, if it is within 2 hours of your next dose, just continue with your regular schedule. Do not double the dose.
- Recent studies have found that for the anti-HIV medications to work, all the medications need to be taken regularly and consistently. Missing or skipping doses of your medication may make it lose its effectiveness as the virus can change itself and become resistant to the medication.

What are the side effects of SAQUINAVIR?

- Common side effects of SAQUINAVIR may include:
 - stomach upset, nausea, gaseousness and diarrhea
 - rash



- headache
- fatigue
- Longer term side effects of SAQUINAVIR may include:
 - increase in the sugar and fat (cholesterol, triglyceride) levels in your blood
 - abnormal body fat distribution (increase in waist and breast size and thinning of the arms and legs and face).
- Consult your doctor or pharmacist if you have these side effects. Do not stop the medication or change the dose before you talk to them.

Can I take SAQUINAVIR with other medications?

- SAQUINAVIR can interact with other drugs. It is important that you tell your doctor and pharmacist about all the prescription and non-prescription medications (including vitamins and herbs) you are taking.
- SAQUINAVIR should not be taken with:
 - Hismanal (astemizole)
 - Seldane (terfenadine)
 - Prepulsid (cisapride)
 - Rifampin
 - Rifabutin

Can I take SAQUINAVIR with alcohol or street drugs?

- It is advisable to avoid excessive amount of alcohol while you are taking SAQUINAVIR. Alcohol may interact with your medications and increase risk of liver damage. Do not skip a dose of your medication because you want a drink.
- SAQUINAVIR may also interact with other street drugs, consult your doctor and pharmacist if you are using street drugs so they can advise you with the necessary precautions.

Can I take SAQUINAVIR if I am pregnant or breastfeeding?

- If you are pregnant and wish to take SAQUINAVIR, please consult your doctor.
- Since the HIV virus can be transmitted through breast milk, breast-feeding is not recommended in HIV positive women.

What other precautions do I need to know when taking SAQUINAVIR?

- You should keep your appointments with your physician for blood tests to check your liver, pancreas and kidney function, blood sugar, cholesterol and triglyceride regularly.
- Make sure you have a continuous supply of the medication.
- SAQUINAVIR does not kill the virus or cure AIDS. It also does not prevent the transmission of HIV, so please remember to always take precautions if you are having sex (e.g. use latex condoms) or using drugs (e.g. use clean syringes).

Copyright @ACAS 2001 & 2003. This fact sheet is produced by ACAS (Asian Community AIDS Services) and is available in English, Chinese (Traditional and Simplified), Tagalog and Vietnamese. Funding for this project is provided by the Ontario HIV Treatment Network (OHTN) and Health Canada. Copies can be downloaded at : www.acas.org/treatment. ACAS provides comprehensive support and case management services for Asian people living with HIV/AIDS; and HIV/AIDS prevention education to the east and southeast Asian Canadian populations in the Greater Toronto Area. ACAS is located at 33 Isabella Street, Suite 107, Toronto, Ontario M4Y 2P7. Tel: (416)-963-4300, Fax: (416)-963-4371 Email: support@acas.org

Legal Disclaimer: While we make every attempt to ensure the accuracy and reliability of information contained in this website/ fact sheet, the information provided here are designed for reference purposes only. These information should not be relied upon as a substitute for medical advice from a qualified professional health care provider and should not be used for diagnosing or treating a condition or illness. Please consult a physician if you have any concerns about your health, treatment regimen and questions related to HIV/AIDS. ACAS, its employees and