



Women's Health

Women and HIV

Last updated September 2003

What symptoms affect women with HIV?

- Women with HIV infection can experience most of the symptoms that men with HIV do. In addition, there are some HIV/AIDS-related conditions that affect only women. These can include menstrual problems, vaginal infections and more frequent diseases of the cervix, such as abnormal growths on the cervix (cervical dysplasia) and cervical cancer.
- Women may also experience different effects from the medications used to treat HIV compared to men. In addition, there are unique concerns about pregnancy and child care that affect HIV positive women.

Menstrual Problems

- Women with HIV may experience changes in their periods. Women can develop irregular cycles, heavier or lighter bleeding or stop menstruating.
- In addition, women with HIV may experience more severe symptoms of PMS (premenstrual syndrome).
- These changes may or may not be directly related to HIV infection. Many other factors, such as long-term stress, severe weight loss, long-term substance use, pregnancy, other chronic illnesses or regular use of some medications can also cause changes in menstrual periods. If you are having any of these problems, they should be checked out by your family doctor and/or a gynecologist (a doctor who specializes in women's reproductive health).

Vaginal Infections

- Vaginal infections, especially yeast infections, are more common in HIV positive women. Because of the weakened immune system, some of these infections may require more aggressive treatment than usual.

Cervical Dysplasia

- Cervical Dysplasia, an abnormal growth or change in the cells of the cervix, occurs more frequently in HIV positive women than HIV negative women. This is a pre-cancerous condition that if untreated, may develop into cervical cancer.
- Fortunately, cervical dysplasia and cervical cancers are preventable. Getting regular pap smears done to screen for these conditions will enable you to get treatment early.

Effects of HIV Drugs

- Using an effective combination of anti-HIV medications can help reduce the amount of virus in your body (your viral load) and improve your body's ability to fight infections by strengthening your immune system. This will often lower the risk of developing infections and symptoms that are related to HIV, including menstrual problems, recurrent vaginal infections and cervical dysplasia.



- On the other hand, women may experience different side effects to some of the HIV drugs than men, and may require different doses. Some women who take protease inhibitors notice changes to their body shape, especially breast enlargement and increase in the size of the waist. In addition, some HIV drugs may affect birth control pills. Please make sure you discuss with your physicians about the effects of all the prescription and non-prescription drugs you are taking, including birth control pills, supplements and herbs.

Pregnancy concerns

- Being HIV Positive does not mean you cannot have children. It also does not mean your baby will definitely be HIV positive. There are treatments that can be taken during pregnancy or at delivery that can reduce the risk of your baby becoming infected by the HIV virus. Please make sure you discuss your concerns with your doctor or midwife.
- HIV can be spread from mother to child through breast milk, so breast-feeding is not recommended in HIV positive women.

Copyright @ACAS 2001 & 2003. This fact sheet is produced by ACAS (Asian Community AIDS Services) and is available in English, Chinese (Traditional and Simplified), Tagalog and Vietnamese. Funding for this project is provided by the Ontario HIV Treatment Network (OHTN) and Health Canada. Copies can be downloaded at : www.acas.org/treatment. ACAS provides comprehensive support and case management services for Asian people living with HIV/AIDS; and HIV/AIDS prevention education to the east and southeast Asian Canadian populations in the Greater Toronto Area. ACAS is located at 33 Isabella Street, Suite 107, Toronto, Ontario M4Y 2P7. Tel: (416)-963-4300, Fax: (416)-963-4371 Email: support@acas.org

Legal Disclaimer: While we make every attempt to ensure the accuracy and reliability of information contained in this website/ fact sheet, the information provided here are designed for reference purposes only. These information should not be relied upon as a substitute for medical advice from a qualified professional health care provider and should not be used for diagnosing or treating a condition or illness. Please consult a physician if you have any concerns about your health, treatment regimen and questions related to HIV/AIDS. ACAS, its employees and board members will not be responsible for any loss or harm, however arising, from the use of, or reliance on this information.